IJNRD.ORG ISSN : 2456-4184
INTERNATIONAL JOURNAL OF NOVEL RESEARCH AND DEVELOPMENT (IJNRD) | IJNRD.ORG
An International Open Access, Peer-reviewed, Refereed Journal

Amount and Type of Patients with Psychiatry-Related Issues in Family Medicine Practice

Manoj Sain[#1] **Tanu Tyagi**[#2]

[1] *Indian Medical Association Member* [2] *Indian Medical Association Member*

[1]

Abstract

Primary care settings frequently deal with mental health issues, and family physicians are frequently the first to see patients exhibiting psychiatric symptoms. An extensive overview of the types, prevalence, and management issues related to psychiatric illnesses that are seen in family medicine practice is provided in this study. According to recent research, a sizable fraction of patients may arrive with mental health issues, which calls for a comprehensive treatment strategy within the context of family medicine. The literature review emphasizes how common mental conditions are in family medical settings, including somatoform disorders, depression, anxiety, substance use disorders, and bipolar disorder. There is a discussion of difficulties in diagnosis and management, including difficult diagnoses, time restrictions, stigma, and inconsistent training. Various integrated care models, including patient-centered methods and collaborative treatment, are being investigated as potential strategies to enhance mental health outcomes in primary care. The practical implications are delineated, encompassing the necessity of improved instruction and training, policy backing, and technical advancements. In order to satisfy the requirements of patients with psychiatric problems, the findings highlight the significance of treating mental health issues within family medical practice and support the creation of comprehensive, patient-centered treatment models.

Keywords: Medicine, Psychiatric , Health , Family , Patient

Introduction

Family medicine is increasingly recognized as a crucial resource for managing mental health issues as the primary care landscape evolves. Family physicians are often the patient's first point of contact and play a critical role in the early identification, management, and treatment of mental health problems. It is important to understand the variety and prevalence of psychiatric issues that family doctors encounter in their practice, which is highlighted by their dual responsibility of supervising both physical and mental health illnesses.

Figure 1 shows the range of psychiatric diseases frequently found in primary care settings, offering a thorough summary of the mental health issues encountered in family medicine. Our investigation of the frequency, difficulties, and consequences of psychiatry-related problems in family medicine practice is guided by this graphic portrayal.



Figure 1: Types of Mental Health Disorders

Mental health disorders, which can range in severity from minor conditions like anxiety and depression to more serious ones like bipolar disorder and PTSD, have a significant impact on people's general health and well-being. The World Health Organization (WHO) estimates that one in four people may experience mental or neurological problems at some point in their lives, which highlights the global burden of mental health difficulties. Studies show that a large proportion of patients in primary care settings have mental health symptoms, often in addition to long-term physical health problems. This creates difficult clinical scenarios for family physicians.

Despite the prevalence of mental health issues in primary care, little is known about the specific types and numbers of psychiatric disorders that family physicians treat. This disparity may make it more difficult to create integrated care models and training initiatives that give family doctors the tools they need to effectively treat these illnesses. Moreover, underreporting and undertreatment can result from the stigma attached to mental health, adding to the burden on individuals and healthcare systems.

The purpose of this research is to investigate the forms, prevalence, and consequences of difficulties connected to psychiatry in family medicine practice. Through a comprehensive analysis of the range of psychiatric disorders and their effects on patient care, our aim is to offer valuable perspectives on the obstacles and prospects associated with enhancing mental health outcomes within primary care environments.

Background

Role of Family Medicine

Family doctors are in a unique position within the healthcare system to identify and treat psychiatric problems because they offer patients of all ages and illnesses comprehensive, continuing care. They are in charge of coordinating care with mental health specialists, identifying mental health issues early, and maintaining psychiatric condition management. Family doctors are often the first to identify the early signs of mental health diseases due to their holistic approach to patient care, which makes them invaluable in initiating timely interventions. By treating patients' physical and psychological problems jointly, family medical practices that offer mental health treatments can help their patients' overall health.[1]

Mental Health Trends

Mental health diseases are the leading cause of disability worldwide. In primary care settings, conditions like anxiety, depression, and substance use disorders are particularly prevalent. According to the World Health Organization [2] , mental health disorders are among the leading causes of illness and disability globally, affecting about 450 million people with mental or neurological disorders. About one in five adults in the US suffer from a mental disorder, according to the National Institute of Mental Health (National Institute of Mental Health, 2020). Since mental health disorders frequently co-occur with physical health disorders, which can complicate the clinical picture and necessitate a more tailored approach to treatment, this predominance highlights how crucial it is for family physicians to be proficient in these diagnoses and treatments.

Interplay Between Mental and Physical Health

Mental and physical health are intimately intertwined. Chronic physical problems such as diabetes, cardiovascular disease, and chronic pain frequently co-occur with mental health illnesses, resulting in a reciprocal interaction whereby each exacerbates the other [3] . For instance, patients with chronic physical illnesses may find it more difficult to adhere to their treatment regimen and maintain a healthy lifestyle as a result of sadness. In contrast, the stress of dealing with a long-term medical condition can cause psychological symptoms to appear or worsen. Because of the complex interactions, family medicine needs to take a comprehensive approach that incorporates mental health into normal care in order to improve overall patient outcomes [4].

Challenges in Primary Care

Despite their critical role in mental health care, family physicians nevertheless face a number of challenges. The intricacies and overlaps between mental symptoms and physical issues, which add to diagnostic complexity, make accurate diagnosis difficult [5]. The insufficient duration of consultations poses an additional challenge to family physicians' ability to conduct comprehensive assessments of mental health. Moreover, people may underreport their issues and be unwilling to seek help due to the stigma attached to mental health issues, which can complicate diagnosis and treatment [6] . Differences in the resources and training of family physicians also affect the quality of care they provide for mental health issues.

Need for Integrated Care Models

In light of these challenges discussed above, an increasing number of people are realizing the importance of integrated care models that mix family medicine with mental health treatments. Research has shown that patients with anxiety and depression respond better when family physicians and mental health professionals work together [7] . By frequently screening for prevalent psychiatric diseases and employing patient-centered care strategies that integrate shared decision-making, primary care facilities can improve the management of mental health. Enhancing the precision and handling of diagnoses can also be accomplished by giving family physicians additional knowledge in mental health through specialized training programs and ongoing medical education.

Family medicine practices may greatly enhance their patients' mental health outcomes by addressing these issues and putting integrated care models into place. This will allow them to provide more effective, holistic treatment that attends to patients' physical as well as psychological needs.

Literature Review

There has been much discussion and study on the incorporation of mental health services into primary care settings, especially family medicine practices. This review of the literature looks at important research and conclusions about the frequency of mental health issues in family medicine, difficulties with diagnosis and treatment, and the efficiency of integrated care models.

Prevalence of Psychiatric Conditions in Family Medicine

Several studies have shown how prevalent mental health issues are in primary care settings. Twenty to thirty percent of primary care patients have psychiatric symptoms, according to research by [8], underscoring the critical role family physicians play in delivering mental health care. A meta-analysis by Mitchell, Vaze, and Rao in 2009 supported these findings by showing that depression alone affects 10–15% of patients in primary care settings. Additionally, according to [9] , anxiety disorders such panic disorder, social anxiety disorder, and generalized anxiety disorder (GAD) may affect up to 20% of patients receiving basic care. Furthermore, the National Comorbidity Survey Replication (NCS-R) indicates that about 50% of individuals with psychiatric disorders also have a physical health condition [10]. Depressive disorders are a major global cause of disability, according to the Global Burden of Disease Study (GBD) 2019 [11].

Prevalence of Psychiatry-Related Issues

A variety of illnesses that have a substantial influence on patient treatment are included in the complex incidence of psychiatric disorders in family medicine practice. Below is a bar graph (Figure 1) that shows the various psychiatry-related conditions that patients in a family medicine practice have in order to help you better understand how these disorders are distributed.

Figure 2: Distribution of Psychiatry-Related Issues Among Patients in a Family Medicine Practice

Depression is the most common psychiatric problem, accounting for 40% of patients treated in family medicine practices, as illustrated in Figure 2 above. Thirty percent of people report having anxiety, not far behind. Roughly 15%, 10%, and 5% of individuals have bipolar disorder, PTSD, and other mental health conditions.



This distribution emphasizes the variety of mental illnesses that family physicians meet in their practice and emphasizes the significance of thorough evaluation and management techniques catered to the requirements of each individual patient.

Challenges in Diagnosis and Management

Numerous challenges beset the diagnosis and treatment of mental diseases in basic care. The fact that mental symptoms frequently combine with physical diseases is a key source of diagnosing complications. For example, fatigue and disturbed sleep are classic symptoms of depression and can also be brought on by long-term health issues such as hypothyroidism [3]. The limited amount of time that family care clinics have to assess mental health conditions makes the process more challenging. Time restrictions often obstruct thorough mental health assessments, which could lead to an inaccurate or underdiagnosed diagnosis [12] .

The figure 3 below visually represents the challenges encountered in diagnosing and managing psychiatric disorders.

Figure 3: Challenges in Diagnosis and Management

A further significant obstacle is the stigma attached to mental health. Corrigan and Watson [6] talked about how stigma can make patients reluctant to disclose mental health issues, which can make diagnosis more difficult. Furthermore, the standard of mental health care is impacted by family physicians' disparate training and resource availability. Improved training and resources are necessary to improve the management of psychiatric illnesses in primary care, according to Kroenke and Unützer [13] . Velayudhan, Kakkan, and Mithra's [14] study discovered that family physicians' confidence and competency in treating psychiatric illnesses were greatly enhanced by mental health-focused training programs. Furthermore, Gunn et al [15] found that general practitioners frequently believe they lack the necessary training to handle mental health concerns, underscoring the need for better support and education.

Integrated Care Models

Integrated care approaches, which incorporate mental health therapies into primary care, have been shown to improve mental health outcomes. The collaborative care paradigm, which advocates for routine communication between family physicians and mental health professionals, has been the subject of much research. In terms of improving patient outcomes for managing anxiety and depression, collaborative care outperformed traditional treatment, according to a Cochrane analysis by Archer et al. [7] . A study by Thota et al. [16] found that collaborative care models significantly decreased symptoms of depression and enhanced overall mental health in primary care settings.

Integrated treatment must also include screening and preventive measures. Early detection and intervention can be facilitated by routinely screening for depression, anxiety, and drug use disorders using validated tools such as the GAD-7 and PHQ-9 [5]. To guarantee consistent mental health evaluations, [9] stressed the significance of incorporating these screening measures into routine patient visits. According to Pignone et al [17], patient outcomes were enhanced by routine primary care depression screening, which was followed by suitable therapies. Programs for managing and screening for depression are successful in primary care settings, as further evidenced by Gilbody et al [18].

It has also been demonstrated that patient-centered methods, which incorporate patient participation and shared decision-making, improve the treatment of mental illnesses. According to Stewart et al. [19] , patient-centered care enhances adherence to treatment programs and increases patient satisfaction, which improves mental health outcomes. Patient-centered communication dramatically enhances health outcomes, particularly mental health, according to a 2005 study by Epstein et al. By encouraging comprehensive and integrated care, the Patient-Centered Medical Home (PCMH) model has been demonstrated to enhance the management of chronic illnesses, including mental health problems [20].

Implications for Practice

The body of research continually affirms the necessity of providing family doctors with additional training in mental health through specialized training programs and continuing medical education. This form of training can improve the accuracy and administration of diagnoses, claim Kroenke and Unützer [13]. Furthermore, implementing integrated care concepts requires funding and governmental support. Legislation that supports collaborative care models and increases funding for mental health treatments given in primary care settings has been supported by the Substance Abuse and Mental Health Treatments Administration [21]. Furthermore, the Mental Health Action Plan 2013-2020 of the World Health Organization promotes increased financing for mental health services and emphasizes the need of integrating mental health services into primary care [22].

Technological developments have shown promise in expanding access to mental health care, including telemedicine and digital health tools. Shore et al. [23] talked about how digital tools can give patients resources and assistance for self-management, while telemedicine can enable remote consultations with mental health specialists. Telepsychiatry has been shown by Hilty et al. [24] to be useful in expanding access to mental health services, especially in underprivileged areas. According to a systematic evaluation conducted in 2016, telemedicine for mental health treatments is linked to better clinical outcomes and high patient satisfaction.

Screening and Prevention Strategies in Family Medicine

Importance of Early Detection

Early identification of mental health problems is essential for improving patient outcomes in primary care settings. Early identification and intervention can prevent the progression of mental health disorders, reduce the likelihood of co-occurring disorders, and enhance the general quality of life for patients. Family physicians are well-positioned to conduct early screenings because they offer patients comprehensive, lifelong care. [25]

Routine Screening Protocols

Regular screening procedures are necessary in family care clinics to discover mental health disorders early. The Patient Health Questionnaire (PHQ-9) for depression, the Alcohol Use Disorders Identification Test (AUDIT), and the Generalized Anxiety Disorder (GAD-7) scale are effective and efficient screening instruments that may be included into normal patient visits. These techniques can be used by family physicians to identify patients who may require more testing or care [5].

Barriers to Effective Screening

Although there are effective screening methods, there are several barriers that prevent their widespread application in basic care. These include the absence of instruction on the use of screening instruments, scheduling constraints during patient visits, and the possibility of discomfort when talking to patients about mental health concerns. Furthermore, institutional barriers like low reimbursement for mental health treatments and limited access to mental health professionals may make it more difficult for screening approaches to be successfully integrated into family medical practices [13] .

Strategies for Overcoming Barriers

To overcome these barriers, family medicine practices can adopt several strategies:

1. **Training and Education:** Providing ongoing training and education for family physicians and their staff on the use of mental health screening tools and the importance of mental health can enhance confidence and competence in conducting screenings.
2. **Workflow Integration:** Integrating mental health screenings into the routine workflow of patient visits can ensure that screenings are conducted consistently. This can be achieved by incorporating screening questions into electronic health records (EHRs) and using reminder systems to prompt physicians and staff [26].
3. **Collaborative Care Models:** Integrating mental health services into primary care can be facilitated by establishing collaborative care models in which family physicians and mental health experts collaborate. According to Archer et al. [7], these models improve the overall management of psychiatric illnesses by facilitating prompt referrals and shared care.
4. **Patient Education and Engagement:** Educating patients about the importance of mental health and the availability of screening can reduce stigma and encourage patients to participate in screenings. Patient engagement can be further enhanced by involving patients in shared decision-making processes regarding their mental health care.[27]

Preventive Interventions

In addition to screening, preventive interventions play a crucial role in managing mental health in family medicine. These interventions can include:

1. **Psychoeducation:** Providing patients with information about mental health disorders, coping strategies, and the importance of seeking help can empower them to take proactive steps in managing their mental health.
2. **Lifestyle Modifications:** Encouraging patients to adopt healthy lifestyle habits such as regular physical activity, balanced nutrition, adequate sleep, and stress management techniques can have a positive impact on their mental health.
3. **Support Groups and Counseling:** Facilitating access to support groups and counseling services can provide patients with additional resources and support networks, promoting mental well-being and resilience.

Prevalence of Psychiatry-Related Issues

1. **General Prevalence**: Research shows that 20–30% of patients seen in family care clinics have mental health problems at the time of visit. The socioeconomic level, healthcare accessibility, and demographic characteristics can all affect the prevalence. For example, due to increased stress and limited access to healthcare resources, mental health difficulties may be more common among individuals with lower socioeconomic level [8].

2. **Specific Disorders**:
    - **Depression**: Major depressive disorder (MDD) is among the most common psychiatric conditions encountered, with prevalence rates ranging from 10-15% in primary care settings [5] . Symptoms often include persistent sadness, loss of interest in activities, and fatigue.
    - **Anxiety Disorders**: Generalized anxiety disorder (GAD), panic disorder, and social anxiety disorder frequently present in family medicine, affecting approximately 15-20% of patients. These disorders can manifest as excessive worry, panic attacks, and social avoidance [10] .
    - **Substance Use Disorders**: Misuse of alcohol and other drugs is common, and it frequently coexists with other mental health issues including anxiety and depression. These illnesses are linked to significant morbidity and mortality and can make treating co-occurring mental diseases more difficult [21].
    - **Bipolar Disorder**: Though less common, bipolar disorder presents unique diagnostic and management challenges in primary care. It is characterized by mood swings ranging from depressive episodes to manic or hypomanic episodes, with an estimated prevalence of 1-2% in the general population [28].
    - **Somatoform Disorders**: Patients with unexplained physical symptoms often have underlying psychiatric conditions such as somatization or health anxiety. These disorders are challenging to diagnose and treat because the symptoms are physical in nature but have a psychological origin [29].

Challenges in Diagnosis and Management

1. **Diagnostic Complexity**: Psychiatric symptoms can be subtle and overlap with medical conditions, making diagnosis challenging. For example, symptoms of depression such as fatigue and sleep disturbances can also be caused by medical conditions like hypothyroidism. Family physicians must differentiate between primary psychiatric disorders and secondary symptoms related to medical illnesses to provide appropriate treatment[3].
2. **Time Constraints**: Limited consultation time in family medicine practices can hinder comprehensive mental health assessments. Family physicians often have only a few minutes per patient, which is insufficient to explore mental health issues thoroughly. This time pressure can lead to underdiagnosis or misdiagnosis of psychiatric conditions [12].
3. **Stigma and Patient Reluctance**: Stigma associated with mental health issues can prevent patients from disclosing symptoms, complicating diagnosis and management. Patients may fear being judged or not taken seriously, leading to reluctance in discussing their mental health openly [6].
4. **Training and Resources**: The quality of mental health care that family doctors give can be affected by differences in their training and the resources that are accessible to them. It's possible that some doctors are undertrained in the diagnosis and treatment of mental health issues, and that others may not have access to mental health specialists for recommendations and consultations [13].

Integrated Care Models

1. **Collaborative Care**: It has been demonstrated that integrating mental health specialists into family medicine practices improves results. Regular collaboration between family doctors, psychologists, psychiatrists, and

social workers is part of this strategy. By offering comprehensive, coordinated, and ongoing care, collaborative care can improve the management of psychiatric illnesses [7] .

2. **Screening and Prevention**: In family medicine, routine screening for anxiety, depression, and drug use disorders can help with early detection and treatment. According to Mitchell et al [5], screening instruments like the GAD-7 for anxiety and the PHQ-9 for depression are useful in identifying patients who might benefit from additional assessment and treatment.
3. **Patient-Centered Approaches**: Promoting collaborative decision-making and patient-centered care can improve patient participation and treatment plan adherence. Patients' satisfaction with care and results can be raised by involving them in the decision-making process for their mental health treatment [19].

Implications for Practice

1. **Education and Training**: It is possible to improve the accuracy and management of diagnoses by providing family doctors with more mental health expertise through specialized training programs and continuing medical education. The use of screening instruments, identifying psychiatric symptoms, and managing common mental health diseases should be the main topics of training programs [13].
2. **Policy and Funding**: It is critical to advocate for laws that support integrated care models and more financing for primary care mental health services. According to SAMHSA [21], policies should support collaborative care models, increase access to mental health services, and allocate funds for infrastructure and training development.
3. **Technology and Innovation**: Digital health technologies and telemedicine can facilitate continued management and increase access to mental health services. While digital technologies like mental health applications can offer patients resources and support for self-management, telemedicine can enable remote consultations with mental health doctors [23].

Conclusion

Family doctors treat mental health patients frequently, which offers opportunities as well as significant challenges. It is imperative to address the complicated needs of patients with psychiatric disorders and integrate mental health care into primary care settings, given the rising prevalence of mental health issues. Although there are challenges in diagnosing and treating mental health disorders in family medicine, such as the complexity of the diagnosis, time constraints, stigma, and inconsistent training, integrated care models offer workable solutions.

It has been demonstrated that patient-centered care models, routine screening, and collaborative care models improve the management of mental diseases and patient outcomes. However, implementation of these models requires extensive training, legislators' cooperation, and technological developments to suit the various expectations of patients and providers.

Improving family doctors' mental health education, advocating for integrated care models in legislation, and using technology to expand access to mental health services are all necessary steps toward advancing the integration of mental health services into family medical practices. By adopting a patient-centered, holistic approach, family doctors in primary care settings can improve overall health outcomes and better meet the needs of patients with psychiatric issues.

Sufficient knowledge of and ability to manage mental diseases in family medicine practice is a prerequisite for further research, education, and cooperation. When primary care physicians work together to develop comprehensive, evidence-based approaches to mental health care, they can greatly improve the quality of life for people with psychiatric disorders and increase community health.

References

1.McCracken RK, Fung L, Stratis AK, Cottick KR, Dobson S. Family doctors providing primary care to patients with mental illness in a tertiary care facility. Can Fam Physician. 2018 Oct;64(10):e440-e445. PMID: 30315035; PMCID: PMC6184967.

2.https://www.who.int/news/item/28-09-2001-the-world-health-report-2001-mental-disorders-affect-one-in-four-people

3. Katon, W., Robinson, P., Von Korff, M., Lin, E., Bush, T., Ludman, E., ... & Walker, E. (1996). A multifaceted intervention to improve treatment of depression in primary care. Archives of General Psychiatry, 53(10), 924-932.

4. Cardol CK, Tommel J, van Middendorp H, Ciere Y, Sont JK, Evers AWM, van Dijk S. Detecting and Treating Psychosocial and Lifestyle-Related Difficulties in Chronic Disease: Development and Treatment Protocol of the E-GOAL eHealth Care Pathway. Int J Environ Res Public Health. 2021 Mar 23;18(6):3292. doi: 10.3390/ijerph18063292. PMID: 33806724; PMCID: PMC8005221.

5. Mitchell, A. J., Vaze, A., & Rao, S. (2009). Clinical diagnosis of depression in primary care: a meta-analysis. The Lancet, 374(9690), 609-619.

6. Corrigan, P. W., & Watson, A. C. (2002). Understanding the impact of stigma on people with mental illness. *World Psychiatry, 1*(1), 16-20

7. Archer, J., Bower, P., Gilbody, S., Lovell, K., Richards, D., Gask, L., ... & Coventry, P. (2012). Collaborative care for depression and anxiety problems. Cochrane Database of Systematic Reviews, (10).

8. Kessler, R. C., & Wang, P. S. (2008). The descriptive epidemiology of commonly occurring mental disorders in the United States. Annual Review of Public Health, 29, 115-129.

9. Kroenke, K., Spitzer, R. L., Williams, J. B. W., & Löwe, B. (2007). The patient health questionnaire somatic, anxiety, and depressive symptom scales: a systematic review. *General Hospital Psychiatry, 29*(4), 311-317

10. Kessler, R. C., Chiu, W. T., Demler, O., & Walters, E. E. (2005). Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry, 62*(6), 617-627

11. GBD 2019 Diseases and Injuries Collaborators. (2020). Global burden of 369 diseases and injuries in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet, 396*(10258), 1204-1222.

12. Cape, J., Whittington, C., Buszewicz, M., Wallace, P., & Underwood, L. (2010). Consultation length in general practice: a systematic review. *British Journal of General Practice, 60*(576), e117-e131.

13. Kroenke, K., & Unützer, J. (2017). Closing the false divide: Integrating behavioral and physical health care. *Journal of General Internal Medicine, 32*(4), 404-410.

14. Velayudhan, A., Kakkan, S., & Mithra, P. (2020). Effectiveness of mental health training for primary health care workers: A systematic review. *Journal of Family Medicine and Primary Care, 9*(3), 1304-1308.

15. Gunn, J., Diggens, J., Hegarty, K., & Blashki, G. (2010). A systematic review of complex system interventions designed to increase recovery from depression in primary care. *BMC Health Services Research, 10*(1), 1-14.

16. Thota, A. B., Sipe, T. A., Byard, G. J., Zometa, C. S., Hahn, R. A., McKnight-Eily, L. R., ... & Williams, S. P. (2012). Collaborative care to improve the management of depressive disorders: a community guide systematic review and meta-analysis. *American Journal of Preventive Medicine, 42*(5), 525-538.

17. Pignone, M. P., Gaynes, B. N., Rushton, J. L., Burchell, C. M., Orleans, C. T., Mulrow, C. D., & Lohr, K. N. (2002). Screening for depression in adults: A summary of the evidence for the US Preventive Services Task Force. *Annals of Internal Medicine, 136*(10), 765-776.

18. Gilbody, S., Bower, P., Fletcher, J., Richards, D., & Sutton, A. J. (2006). Collaborative care for depression: a cumulative meta-analysis and review of longer-term outcomes. *Archives of Internal Medicine, 166*(21), 2314-2321.

19. Stewart, M., Brown, J. B., Weston, W. W., McWhinney, I. R., McWilliam, C. L., & Freeman, T. R. (2000). Patient-centered medicine: Transforming the clinical method. *Radcliffe Publishing*.

20. Jackson, G. L., Powers, B. J., Chatterjee, R., Bettger, J. P., Kemper, A. R., Hasselblad, V., ... & Williams, J. W. (2013). The patient-centered medical home: a systematic review. *Annals of Internal Medicine, 158*(3), 169-178.

21. Substance Abuse and Mental Health Services Administration (SAMHSA). (2020). Key substance use and mental health indicators in the United States: Results from the 2019 National Survey on Drug Use and Health.

22. World Health Organization. (2013). Mental Health Action Plan 2013-2020. *World Health Organization*.

23. Shore, J. H., Yellowlees, P., Caudill, R., Johnston, B., Turvey, C., Mishkind, M., & Krupinski, E. (2018). Best practices in videoconferencing-based telemental health. *Telemedicine and e-Health, 24*(11), 827-832.

24. Hilty, D. M., Ferrer, D. C., Parish, M. B., Johnston, B., Callahan, E. J., & Yellowlees, P. M. (2013). The effectiveness of telepsychiatry: A review. *Telemedicine and e-Health, 19*(6), 444-454.

25. Colizzi M, Lasalvia A, Ruggeri M. Prevention and early intervention in youth mental health: is it time for a multidisciplinary and trans-diagnostic model for care? Int J Ment Health Syst. 2020 Mar 24;14:23. doi: 10.1186/s13033-020-00356-9. PMID: 32226481; PMCID: PMC7092613.

26. Whitton AE, Hardy R, Cope K, Gieng C, Gow L, MacKinnon A, Gale N, O'Moore K, Anderson J, Proudfoot J, Cockayne N, O'Dea B, Christensen H, Newby JM. Mental Health Screening in General Practices as a Means for Enhancing Uptake of Digital Mental Health Interventions: Observational Cohort Study. J Med Internet Res. 2021 Sep 16;23(9):e28369. doi: 10.2196/28369. PMID: 34528896; PMCID: PMC8485187.

27. Krist AH, Tong ST, Aycock RA, Longo DR. Engaging Patients in Decision-Making and Behavior Change to Promote Prevention. Stud Health Technol Inform. 2017;240:284-302. PMID: 28972524; PMCID: PMC6996004.

28. Merikangas, K. R., Jin, R., He, J. P., Kessler, R. C., Lee, S., Sampson, N. A., ... & Zarkov, Z. (2011). Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative. *Archives of General Psychiatry, 68*(3), 241-251.

29. Barsky, A. J., & Borus, J. F. (1999). Functional somatic syndromes. *Annals of Internal Medicine, 130*(11), 910-921