



PHARMACOGNOSTIC PRESPECTIVE OF *Artocarpus heterophyllus Lam* LEAF AND ITS INVITRO ANTIRHEUMATIC ACTIVITY

¹SHONE SANTHOSH, ²SIYANA. N, ³SAHANA S.S

¹Student, Ezhuthachan College Of Pharmaceutical Sciences, Marayamuttom, Neyyattinkara,
Trivandrum, Kerala, India, 695124.

²Student, Ezhuthachan College Of Pharmaceutical Sciences, Marayamuttom, Neyyattinkara,
Trivandrum, Kerala, India, 695124.

³Student, Ezhuthachan College Of Pharmaceutical Sciences, Marayamuttom, Neyyattinkara,
Trivandrum, Kerala, 695124

ABSTRACT

Artocarpus heterophyllus Lam, popularly known as Ceylon jack tree or jackfruit, evergreen tree reaching 8-25 m high, belongs to the family Moraceae. The trunk is straight, rarely buttressed, with a circumference of 50-100 cm. The stems are straight, forming a 30° angle with the trunk. The leaves are alternate, simple, pinnately veined, coriaceous, glossy, 4-25 cm long x 2-12 cm, dark green on the upper side and pale green beneath. The leaf has many therapeutic properties especially its diabetes control, antioxidant and anti-aging properties. Traditionally leaves are useful in fever, boils, wounds and skin diseases. In the present study, we discuss and highlight the recent findings concerning the different classes of RA therapies including the conventional and modern drug therapies. Approximately, one per cent of the world population lives with rheumatoid arthritis (RA), with at least twice as many women affected as men. Selection of leaves are important as compared to the other parts because it includes more constituents and also can be cheaply available.

KEYWORDS: *Artocarpus heterophyllus Lam*, Rheumatoid arthritis, Inflammation, Moraceae

INTRODUCTION

Medicinal plants provide major source of molecules with medicinal properties due to presence of natural compounds. Medicinal plants are useful for curing human diseases and play an important role in healing due to presence of phytochemical constituents. Plant is an important source of medicine and plays a key role in world health. Medicinal herbs or plants have been known to be an important potential source of therapeutics or curative aids. The use of medicinal plants has attained a commanding role in health system all over the world. This involves the use of medicinal plants not only for the treatment of diseases but also as potential material for maintaining good health and conditions. Chemically prepared drugs may act quickly, but they have side effects which affect human body negatively in the long run, whereas, medicinal plants work in an integrated or pro-biotic with little or no adverse effects on the body.

Chronic diseases—including, cancer, diabetes, hypertension, stroke, heart disease, respiratory diseases, arthritis, obesity, and oral diseases—can lead to hospitalization, long-term disability, reduced quality of life, and death. Chronic diseases are characterized by high prevalence among populations, rising complication rates, and increased

incidence of people with multiple chronic conditions. Approximately, one per cent of the world population lives with rheumatoid arthritis (RA), with at least twice as many women affected as men. The World Health Organization estimates more than 23 million people live with RA. Rheumatoid arthritis (RA) is one of more than 100 types of arthritis. This chronic autoimmune disorder affects the lining of synovial joints in about 0.5% of people and may induce severe joints deformity and disability. Rheumatoid arthritis (RA) is one of more than 100 types of arthritis. This chronic autoimmune disorder affects the lining of synovial joints in about 0.5% of people and may induce severe joints deformity and disability.

Rheumatoid arthritis impacts health life of people from all sexes and ages with more prevalence in elderly and women people. Non-steroidal anti inflammatory drugs (NSAIDs) (e.g.; Naproxen, Aspirin, Ibuprofen, oxaprozin, salsalate, diclofenac, mefenamic acid, meclofenamate, ketorolac etc.), corticosteroids (e.g.; Prednisone, Methylprednisolone, dexamethasone etc.), and disease-modifying antirheumatic drugs (DMARDs) (e.g.; Methotrexate, Hydroxychloroquine, sulfasalazine etc.) remain the most known treatments used against rheumatoid arthritis. Side effects such as Gastric irritation, Peptic ulcer, Gastrointestinal bleeding, Nausea, Abdominal discomfort, Kidney impairment, Cardiovascular effects, Rash, Diabetes, Osteoporosis, Metabolic syndrome, Hepatic dysregulations, Pneumonitis, Hematologic disorders, Infections, Nephrotoxicity, Hypercholesterolemia, Creatininemia, cellulitis, diarrhea, headache, hypertension, liver failure etc. (Reda Ben Mrid et al 2022).

However, not all patients respond well to these drugs and therefore, new solutions are of immense need to improve the disease outcomes. In the present review, we discuss and highlight the recent findings concerning the different classes of RA therapies including the conventional and modern drug therapies. Traditional medicinal plants are practiced worldwide for treatment of arthritis especially in developing countries where resources are meager. 485 plant species belonging to 100 families, traditionally used in arthritis. Several herbal remedies are promoted today for treating arthritis, including turmeric, ginger, *Boswellia serrata*, devil's claw, willow bark extract and feverfew etc. (Manjusha Choudhary 2015).

Artocarpus genus (Family: Moraceae-mulberry family) received a great level of scientific interest as they consist of therapeutically active secondary metabolites and is economic source of food and widely used in traditional medicine. *Artocarpus* species are used as food and for traditional folk medicine in South-East Asia, Indonesia, Western part of Java and India. *Artocarpus* plants offer advantages as a profitable multipurpose crop for producing fruits and timber. *Artocarpus* has long been recognized and economically is of appreciable importance as a source of edible aggregate fruit; such as *Artocarpus heterophyllus* (Jack fruit), *Artocarpus altilis* (bread fruit) and *Artocarpus chempeden* (Chempedak) and yielding fairly good timber. *Artocarpus heterophyllus* popularly known as jack fruit is one of the important and commonly found trees in the home gardens of India and Bangladesh. The term jackfruit is derived from the Portuguese word Jaca which in turn is adopted from the word “Chakka” of Malayalam - A regional Indian language. Extracts of its plant parts have been applied in traditional medicine for the treatment of diarrhea, diabetes, malarial fever, tapeworm infestation, and as wound healing, antisyphilitic, expectorant and also to treat anaemia, asthma and dermatitis.

KINGDOM	PLANTAE
CLADE	TRACHEOPHYTES
CLADE	ANGIOSPERMS
CLADE	EUDICOTS
CLADE	ROSIDS
FAMILY	MORACEAE
GENUS	ARTOCARPUS
SPECIES	A.heterophyllus

Taxonomy of *Artocarpus heterophyllus lam*

Leaves are thin-leathery and obovate-elliptic to elliptic. The leaves are smooth on the surface and rough on the undersides. Stipules are ovate-acute, deciduous and leave ring-like form on the twigs. The leaf has many therapeutic properties especially its diabetes control, antioxidant and anti-aging properties. It has distinctive flavour and fragrance and it is known be rich in vitamin. Jackfruit leaves has several therapeutic effect and work wonders when taken (especially on an empty stomach). Wide variations in leaf blade shape i.e., elliptic, obovate, oblong, broadly elliptic and narrowly elliptic were recorded in the selected jackfruits. Different fruit shapes like ellipsoid, spheroid, oblong, clavate, oblong and irregular were recorded in the selected accessions. Flake texture and flake flesh colour also showed wide variation. In seeds, different seed shapes such as ellipsoid, irregular, reniform, spheroid and oblong were recorded.

A. heterophyllus Lam. (AH), popularly known as Ceylon jack tree or jackfruit, is considered to be “poor man’s food”, as it is widely available in summer at an economical price when there is a shortage of agricultural produce in India. In India, it is also called “Panasa”, “Atibruhatphala”, “Kantaphal”, and “Kanthal” due to its physical characteristics. The *A. heterophyllus* tree is also considered to be a source of flavonoids, sterols, volatile acids, carotenoids, and tannins, which vary depending on the stage of maturity. Moreover, it also has a unique combination of iron, vitamin C, vitamin B complex (especially vitamin B6, niacin, folic acid, and riboflavin), and minerals (mainly calcium and potassium). The fruit of *A. heterophyllus* has been employed in value-added products such as fruit juice, dried chips, jam, candies, jelly, marmalades, leather, and ice cream, depending on the sensorial characteristics at different stages. *A. heterophyllus* bears flowers after 6–8 years of planting, and flowers start to bloom from November to March. It is reported that, being a monoecious tree, the flowers are covered by two spathes of 5–10 cm in length. The central and peripheral regions bear male flowers, while footstalks from the main trunk bear female flowers (generally round in shape). The distinction between the flowers can be made based on surface characteristics: female flowers are generally large with a rough surface, whereas male flowers are small with a smooth surface. The leaves and stem show the presence of sapogenins, cycloartenone, cycloartenol, β sitosterol and tannins, they show estrogenic activity. A root contains β -sitosterol, ursolic acid, Betulinic acid and cycloartenone.

Traditionally leaves are useful in fever, boils, wounds and skin diseases. The young fruits are acrid, astringent, and carminative. The ripe fruits are sweet, cooling, laxative, aphrodisiac and also used as a brain tonic. The seeds are, diuretic, and constipating. The wood is nervine, antidiabetic, sedative and is useful in convulsions. The latex is useful in dysopia, ophthalmic disorders and pharyngitis and also used as antibacterial agent. The ash of Jackfruit leaves is used in case of ulcers. The root is a remedy for skin diseases and asthma. An extract of the root is taken in cases of fever and diarrhea. The bark is made into poultices. Heated leaves are placed on wounds. The wood has a sedative property and its pith is said to be abortifacient. Latex is used as an anti-inflammatory agent.

Nature has blessed us with enormous wealth of herbal plants which are widely distributed all over the world as a source of therapeutic agents for the prevention and cure of various diseases In India, there are more than 2500 plants species which are currently used as herbal medicaments. For than 3000 years, the herbal medicines are used either directly as folk medication or indirectly in the preparation of recent pharmaceuticals. Selection of leaves are important as compared to the other parts because it includes more constituents and also can be cheaply available. In India we don't give more importance to the *A. heterophyllus* leaf. Usually, they are burning along with other waste products. So, we will select to make an advantage from that.

MATERIAL AND METHOD

COLLECTION OF SAMPLES AND AUTHENTICATION

Fresh Leaf part of *Artocarpus heterophyllus* belonging to the family Moraceae was collected from kazhakkotam city at Trivandrum district of kerala. The identification and authentication of the medicinal plant were carried out by Dr.E.A. Siril Proffesor and head, Department of Botany, University of Kerala, Kariavattom, Thiruvananthapuram. The drug was shade dried, stored in airtight container and labelled. The drug was powdered and used for extraction, pharmacognostical, physico-chemical parameters, phytochemical, and pharmacological study. The RBC cell was collected from a laboratory in Trivandrum.

PHARMACOGNOSTICAL STUDY

The Leaf sample was subjected to morphological and microscopical examination

MORPHOLOGY STUDY

The leaf part of plant *A. heterophyllum* Lam was collected and subjected to morphological studies such as shape, size, colour, odour, taste.

MICROSCOPY STUDY

The sample was subjected to transverse section. After collecting the plant material, cut the material with razor blade to a thickness of 10- 20 μm and examined the sample under a microscope after phloroglucinol and HCL.

POWDER MICROSCOPY

Spread a small quantity of the powder on the slide, treated with acetic acid and observed under microscope.

QUANTITATIVE LEAF MICROSCOPY

VEIN ISLET NUMBER

Vein – islet number is the minute area of photosynthesis tissue encircled by the ultimate division of the conducting strands. A piece of the leaf was cleared by boiling in chloral hydrate solution and camera lucida and drawings board were arranged and 1 mm line was drawn with help of stage mm. A square was constructed on this line in the centre of the field. The slide was placed on the stage. The veins included within the square were traced off, completing the outline of those islets which overlap two adjacent side of the square. The average number of vein islet from our adjoining square, to get the value for one square mm was calculated.

VEIN TERMINAL NUMBER

Vein termination number is the number of veinlet terminations per mm of leaf surface. A piece of the leaf was cleared by boiling in chloral hydrate solution and camera lucida and drawings board were arranged and 1 mm line was drawn with help of stage mm. A square was constructed on this line in the centre of the field. The slide was placed on the stage. The veins included within the square were traced off, completing the outline of those islets which overlap two adjacent side of the square. The number of veinlet termination present within the square was counted and the average number of veinlet termination number from the four adjoining square to get the value for 1 square mm was found known as vein termination number.

STOMATAL NUMBER

Stomatal number is the average number of stomata per square millimeter of epidermis. A piece of leaf was cleaned and the upper and lower epidermis was peeled out separately by means of forceps. It was kept on slide and mounted in glycerin water. Camera lucida was attached and drawing board was placed for drawing the cells. A square of 1 mm by means of stage micrometer was drawn on it. The slide with cleared leaf was placed on the stage and the epidermal cells and stomata were traced. The number of stomata and the number of epidermal cells in each field were counted.

STOMATAL INDEX

The percentage proportion of the ultimate divisions of the epidermis of a leaf which can be converted into stomata is termed as stomatal index. A piece of leaf was cleaned and the upper and lower epidermis was peeled out separately by means of forceps. It was kept on slide and mounted in glycerin water. Camera lucida was attached and drawing board was placed for drawing the cells. A square of 1 mm by means of stage micrometer was drawn on it. The slide with cleared leaf was placed on the stage and the epidermal cells and stomata were traced. The number of stomata and the number of epidermal cells in each field were counted. The numbers of stomata were counted as stomatal number and the stomatal index using the above formula was calculated separately for upper and lower surface. Stomatal index can be calculated by using following equation:

$$I = S / E + S \times 100$$

where,

I = stomatal index.

S = number of stomata per mm².

E = number of ordinary epidermal cells per mm².

PHYSICAL STUDY

ASH VALUE

Ash content represents the inorganic residue remaining after ignition and complete oxidation of organic material. Ash values are helpful in determining the quality and purity of crude drug especially in powdered form.

PROCEDURE:

Total Ash

Weighed and ignited flat, thin teared silica crucible and weighed about 2gm of powdered plant material into the crucible. Heated with a burner using a flame about 2cm height and supporting the dish about 7cm above the flame. Heated till vapour almost cease to be evolved, then lower the dish and the heat moved strongly until all the carbon is burned off, cooled in a desicator. Weighed the ash and calculate the total ash.

Acid – insoluble Ash

Proceed as per the step mentioned in the procedure for total ash value of crude drug. Further using 25 ml of dil.HCL washed the total ash from the dish used for the total ash into 100 ml beaker. Placed wire gauze over a burner and boiled for 5 min. Filtered through as ashless filter paper, washed the residue twice with hot water. Ignited a crucible in a flame cooled and weighed. Put the filter paper and residue together into the crucible, heated gently until vapour cease to the evolved and then moved strongly until carbon has been released. Cooled in a desicator and weighed the residue and calculated acid – insoluble ash value of the drug.

Water – Soluble Ash

Using 25 ml of water, washed the total ash from the dish used for the total ash into 100ml beaker. Placed wire gauze over the burner and boiled for 5 min. Filtered through as ashless filter paper, washed the residue twice with hot water. Ignited the crucible in a flame cooled and weighed. Put the filter paper and residue together in the crucible, heated gently until vapour ceases to the evolved and then moved strongly until carbon has been released. Cooled in a desicator, weighed the residue and calculate acid – soluble ash of the drug.

LOSS ON DRYING

Moisture content was determined by loss of water in terms of percent W/W by using the formula: Moisture content = Fresh weight – Dry weight / Fresh weight x 100. Procedure: Weighed an empty Petri dish and taken 5gm of sample into it. Put this sample in an oven and adjusted 105°C temperature and this sample in an oven for 3 hrs. Cooled the petri dish in a desicator and weighed the loss in weight is usually recorded as moisture.

EXTRACTIVE VALUE

Extractive value helps to determine the number of active constituents present in the plant materials. Procedure:

I. Alcohol soluble extractive value

Weighed accurately about 5 gm of powdered plant material into 250 ml conical flask with stopper and added 100 ml of alcohol. Shaken the flask frequently during first 6 hrs. Kept it aside without disturbing for 18 hrs and then filtered. Pippeted out 25 ml of filterate and evaporated to dryness in a weighed shallow flat-bottomed dish on a

water bath. Then dried the residue at 105 °C to a constant weigh and calculated the percentage of alcohol soluble extractive value.

$$\% \text{ of alcohol soluble extractive value} = \text{weight of residue} / \text{weight of drug} \times 100$$

II. Water soluble extractive value

Weighed accurately about 5 gm of powdered plant material into 250 ml conical flask with stopper and added 100 ml of chloroform water. Shaken the flask frequently during first 6 hrs. Kept it aside without disturbing for 18 hrs and then filtered. Pippeted out 25 ml of filtrate and evaporated to dryness in a weighed shallow flat-bottomed dish on a water bath. Then dried the residue at 105 °C to a constant weigh and calculated the percentage of water- soluble extractive value.

$$\% \text{ of water-soluble extractive value} = \text{weight of residue} / \text{weight of drug} \times 100$$

PHYTO CHEMICAL STUDY

Extracts obtained were subjected to various chemical test to the chemical constituents presents in them. Results were tabulated.

TEST FOR ALKALOIDS

The test solution was prepared by dissolving the extract in dilute HCL, the solution was filtered. The filtrate was then subjected to the following test for the detection of the presence of alkaloids.

1. Dragendroff's test: To 2ml of the filtrate was added to 1 ml of dragendroff's reagent (solution of potassium bismuth iodide). Formation of orange or reddish – brown precipitate indicates the test as positive.
2. Mayer's test: To 1 ml of test solution or filtrate was added a drop or two of the Mayer's reagent (Potassium mercuric iodide.) Formation of yellow cream precipitate indicates the presence of alkaloids.
3. Hager's test: To 1 ml of the filtrate or solution, added one drop or two drop of Hager's reagent (saturated picric acid solution). Formation of yellow precipitate indicated the presence of alkaloids.
4. Wagner's test: Two drops of Wagner's reagent (Iodine in potassium iodide). Formation of yellow or brown precipitate indicate the presence of alkaloids.

TEST FOR CARBOHYDRATES

The test solution was prepared by dissolving the extract individually in 5ml distilled water and filtered. The filtrates were used to test for the presence of carbohydrates.

1. Molisch's test: Filtrates were treated with 2 drops of alcoholic α -naphthol in a test tube and 2ml of con. sulphuric acid was added carefully along the sided of the test tube. Formation of violet ring at the junction indicates the presence of carbohydrates.
2. Benedict's test: Filtrates were treated with Benedict's reagent and heated on water bath. Formation of orange red precipitate indicates the presence of carbohydrates.
3. Fehling's test: Filtrates were hydrolyzed with diluted hydrochloric acid, neutralized with alkali and heated with Fehling's A and B solutions. Formation of red precipitate indicates the presence of reducing sugars.

TEST FOR GLYCOSIDES

Extracts were hydrolyzed with diluted hydrochloride acid and then subjected to test for glycosides.

1. Keller -killani test: Filtrate were treated with glacial acetic acid containing a drop of ferric chloride. Formation of brown coloured ring indicates the presence of glycosides.

TEST FOR PROTEINS

Extracts were prepared in water with phosphate buffer, soluble once are easily extracted by distilled water and insoluble once are dissolved by alkaline treatment, then subjected to test for protein.

1. Biuret test: Filtrate were added with 1ml of 4 % NaOH solution and a few drops of 1% copper sulphate solution. Formation of violet or pink color indicates the presence of protein.

2. Million's test: Filtrate were treated with million's reagent. Formation of white precipitate, on warming the precipitate of brick red or it may be dissolved indicates the presence of protein.

TEST FOR AMINO ACID

Extracts was dissolved in water and subjected to test for amino acid.

1. Ninhydrin test: Filtrate were treated with few drops of Ninhydrin solution and solution is treated in water bath for 10 minutes. Formation of purple or bluish colour indicates the presence of amino acid.

TEST FOR SAPONINS

Extracts were dissolved in distilled water and subjected to test for saponins.

1. Foam test: Filtrate were placed into a test tube and shaken vigorously for 2 minutes. Formation of persistent appearance of foam lasting for an at least 15 minutes indicates the presence of saponins.

TEST FOR FLAVONOIDS

Extracts were treated with 2M HCL and heated for about 30-40 min at 100°C. The extract was cooled and again extracted with ethyl acetate which was further concentrated to dryness and subjected to test for flavonoids.

1. Shinoda's test: Filtrate were treated with 95% v/v methanol and few drops of Con.Hcl and 0.5 gm magnesium turnings. Formation of pink color indicates the presence of flavonoids.

2. Lead acetate test: Filtrate were treated with equal amount of lead acetate solution. Formation of yellow-coloured precipitates indicates the presence of flavonoids.

3. Sodium hydroxide test: Filtrate were treated with large amount of NaOH. Formation of yellow colour which decolorized after the addition of acid indicator.

TEST FOR TANNINS

Extract was dissolved in water and alcohol which is subjected to test for tannins.

1. Ferric chloride test: Filtrate were treated with 1ml of ferric chloride. Formation of dark blue or greenish black indicates the presence of tannins.

2. Gelatin test: Filtrate were treated with few ml of 1% gelatin solution containing 10% NaCl. Formation of white principles indicates the presence of tannins.

3. Lead acetate test: Filtrate were treated with few ml of 10% lead acetate solution. formation of white precipitate indicates the presence of tannins.

TEST FOR STEROIDS AND TRITERPENOIDS

Extract was dissolved in chloroform and subjected to test

1. Salkowski test: Filtrate were treated with 2ml chloroform and 3ml concentrated sulphuric acid carefully added to form a layer. Formation of reddish-brown coloration of the interface indicates the presence of terpenoids.

2. Liebermann-Burchard test: Filtrate were filled with chloroform, acetic anhydride and few drops of Con.sulphuric acid. Formation of dark green color indicates the presence of terpenoids.

TEST FOR FATS AND OILS

1. Filter paper test: Filter paper gets permanently stained with oils.

VOLATILE OIL EXTRACTION

Laboratories make use of the Clevenger apparatus for this purpose. The name of this apparatus was derived from its inventor, Joseph Franklin Clevenger in 1928. The Clevenger apparatus has three main parts- a round bottom flask in which the organic material is placed, a separator for automatically separating the distilled solution and a condenser. The apparatus is available in varied sizes to facilitate organic material. It looks similar to a glass double boiler having several valves to adjust the temperature. The process of extracting oil from an organic source involves the use of steam. The steam helps in avoiding the degradation of the essential oils. At laboratory scale, we bring a few liters of water to a boil, and steam rises in a column containing the more or less finely ground plant. The vapor phase is then directed to a condenser, and the liquid is collected in a graduated burette. Thanks to a bended connection at the base of the burette, the hydrosol flows to the left in the beaker, while the essential oil remains in the burette. After typically 2 h of extraction (calculated from the first condensed drop), we can measure the volume of oil recovered and calculate the yield from the mass of plant introduced. This technique has the advantage of allowing us to recover the hydrosol. It also allows us to perform extractions on larger quantities of plants.

TOXICITY STUDIES

PRELIMINARY TOXICOLOGICAL SCREENING USING DAPHNIA MAGNA

Select a range of concentrations that span those causing zero mortality to those causing complete mortality.

Rinse all exposure chambers, except the chamber containing 100% test media, in dilution water. Mix concentrations and pour into each exposure chamber and measure 0.5 mL of the test media into a beaker and dilute to 500 ml. Using a graduated cylinder, pour out 50 mL into each exposure chamber and pour the rest into a beaker for chemical measurements. Continue these steps for all concentrations. Always work from the lowest concentration to the highest in order to minimize the risk of cross contamination. Using a wide bore pipette, randomly select and carefully place daphnia into each exposure chamber. Place the pipette tip below the surface and gently expel each daphnia individually into the chamber. The test begins when half of the organisms are in the exposure chambers. Measure and record mortality and survival at one hour and then at 24 and 48 hr. Measure and record temperature, dissolved oxygen, pH, conductivity, alkalinity, and hardness after the test begins and at the completion of the test. The test is complete at the end of 48 hours.

PHARMACOLOGICAL STUDY

INVITRO HRBC MEMBRANE STABILIZATION

HRBC method was used for the estimation of anti-inflammatory activity in vitro. The RBC cells was collected and washed with isosaline solution and a 10% v/v suspension was made with isosaline. This HRBC suspension was used for the estimation of anti-inflammatory property. Different concentrations of extract, reference sample and control were separately mixed with 1mL of phosphate buffer, 2 mL of hyposaline and 0.5 mL of HRBC suspension. All the assay mixtures were incubated at 37 °C for 30 minutes and centrifuged at 3 000 rpm. The supernatant liquid was decanted and the hemoglobin content was estimated by a spectrophotometer at 560 nm. The percentage hemolysis was estimated by assuming the hemolysis produced in the control as 100%.

$$\text{Percentage protection} = 100 - (\text{OD sample} / \text{OD control}) \times 100$$

Diclofenac sodium(5mg/ml) was chosen as the standard drug. The hemoglobin content in the supernatant solution and standard drug solution was estimated by using spectrophotometer at 560 nm.

RESULT AND DISCUSSION

MORPHOLOGICAL STUDY

Shape: Alternate, entire, simple, glossy, leathery, stiff, large and elliptic to oval in form with an entire margin, blunt apex, tapers to base, glabrous surface, Netted pinnate venation. Leaves are often deeply lobed when juvenile and on young shoots. Size: 8 -10 cm in width, 15-17cm length. Colour: Green colour. Odour: Characteristic odour. Taste: Bitter taste.

MICROSCOPIC STUDY

TS of the leaf shows a dorsiventral structure with slightly wavy upper and much wavier lower epidermis. Mesophyll shows the presence of a single layer of compacted elongated palisade cells which is followed by spongy parenchyma. It consists of cambium, phloem, ground tissue, pith, xylem, vascular bundles.

Powder microscopy showed xylem vessels, epidermal cells with anomocytic stomata, covering trichomes, rosette form of calcium oxalate crystals and starch grains.

QUANTITATIVE MICROSCOPY

Quantitative microscopy of leaf constants

SL.NO	QUANTITATIVE MICROSCOPY	OBSERVATION
1	STOMATAL NUMBER	15.6
2	STOMATAL INDEX	25.7
3	VEIN ISLET NUMBER	15.4
4	VEIN TERMINATION NUMBER	30.7

Table : Quantity microscopy of leaf of *A.heterophyllus.Lam* leaf

The leaf constants of *A.heterophyllus.Lam* was examined and found to be the following observations.

Stomatal number:15.6

Stomatal index:25.7

Vein-islet number:15.4

Vein termination number:30.7

PHYSICAL STUDY

Parameters	Leaf Values % W/W
Water- soluble extractive value (% W/W)	5.4
Alcohol- soluble extractive value (%W/W)	6.0
Loss on drying at 105°C (%W/W)	0.168
Total ash value (% W/W)	27.0
Acid-insoluble ash value (% W/W)	14.07
Water-soluble ash value (% W/W)	24.80

Physical study of *A.heterophyllus.Lam* leaf

The physical study of *A. heterophyllus.Lam* leaf was carried out and found the values of water extractive values, alcohol extractive values, total ash, water soluble ash, acid insoluble ash and loss on drying.

PHYTO CHEMICAL STUDY

SL.NO	NAME OF TEST	ALCOHOL EXTRACT	WATER EXTRACT
1	Carbohydrate	+ve	+ve
2	Alkaloid	+ve	-ve
3	Glycoside	+ve	-ve
4	Flavonoids	+ve	+ve
5	Tannins	-ve	+ve
6	Protein	+ve	+ve
7	Amino Acid	+ve	+ve
8	Saponin	+ve	+ve
9	Steroids And Triterpenoids	+ve	+ve
10	Fats And Oils	-ve	-ve

Table : Phyto chemical study of *A.heterophyllus.Lam* leaf

The phytochemical study was carried out and the ethanolic extract of *A.heterophyllus.Lam* contain the presence of carbohydrate, alkaloid, glycoside, flavanoids, protein, saponin, steroids and triterpenoids. The water extract contains the presence of tannins, protein, amino acid. saponin, steroid and triterpenoids.

Volatile oil extraction method was carried out using Clavancher apparatus and found the absence of oil in *A.heterophyllus.Lam* leaf part.

TOXICITY STUDIES

PRELIMINARY TOXICOLOGICAL SCREENING USING DAPHNIA MAGNA

The methods used to determine the LC50 differ depending on the results of the test. If there is no partial mortality in any replicate (i.e., all alive or all dead), then the Moving-Average Method may be used to determine the LC50. If there is partial mortality within a replicate, then the Probit Method should be used to calculate the LC50. Also, the Lowest Observable Effect Concentration (LOEC) is recorded and the No Observable Effects Concentration (NOEC) is recorded. Since this is a simple acute test, only mortality is recorded upto 2000µg/ml and no mortality was observed

TEST CONCENTRATIONS(µg/ml)	MORTALITY
100	Nil
500	Nil
1000	Nil
1500	Nil
2000	Nil

Table : Toxicity study of *Daphnia magna*

PHARMACOLOGICAL STUDY INVITRO HRBC MEMBRANE STABILIZATION

HRBC membrane stabilization was carried out after collecting RBC cells from the laboratory. Then alcohol extract of *A.heterophyllus.Lam* was added to the HRBC at different dilution (50,100 & 150 µg/ml) and then added phosphate buffer and hyposaline solution and it was again centrifugated at 3000 rpm and the solution was incubated at 37°C for 30 minutes, then analysed at 560 nm. The percentage haemolysis was estimated by assuming the hemolysis produced in the control as 100%. Diclofenac sodium was used as the standard drug (5mg/ml).

	ABSORBANCE (nm)
CONTROL	0.854
DICLOFENAC SODIUM (STANDARD)	0.686
ETHANOLIC LEAF EXTRACT CONCENTRATION	
50 µg/ml	0.45
100 µg/ml	0.52
150 µg/ml	0.60

Table :Pharmacological study of *A.heterophyllus.Lam*

ANTI RHEUMATIC ACTIVITY IN HRBC USING ETHANOL LEAF EXTRACT




Fig : Graphical representation of anti rheumatic activity in HRBC using ethanolic leaf extract

A. heterophyllus.Lam ethanolic extracts at different concentrations (50,100,150 µg/mL) showed significant stabilization towards HRBC membranes. The percentage protection of ethanolic extract at concentration 150 µg/mL was higher than that of other concentrations. However, the percentage protection was found to be increased at higher concentration.

CONCENTRATIONS	PERCENTAGE
	(%)
STANDARD	39.7
50 µg/ml	10.5
100 µg/ml	15.12
150 µg/ml	25.74

REFERENCE

1. Ahasan Ullah Khan, Israt Jahan Ema, Md. Ruman Faruk, Shofiul Azam Tarapder, Anayat Ullah Khan, Sana Noreen and Muhammad Adnan. A Review on Importance of *Artocarpus heterophyllus* L. (Jackfruit): Journal of Multidisciplinary Applied Natural Science Vol. 1 No. 2 (2021).
2. Alamgeer Umme Habiba Hasan. Study of in vitro and in vivo therapeutic efficacy of *Ribes alpestre* Decne in Rheumatoid arthritis: Brazilian journal of pharmaceutical science 2019.
3. Baliga MS, Shivashankara AR, Raghavendra Haniadka c, Jerome Dsouza c, Harshith P. Bhat d. Phytochemistry, nutritional and pharmacological properties of *Artocarpus heterophyllus* Lam (jackfruit): A review. Food Research International 2011; 44:18001811.
4. Chackrewarthy S, Thabrew MI, M. K. B. Weerasuriya, and S. Jayasekera. Study of the hypoglycemic and hypolipidemic effects of an ethylacetate fraction of *Artocarpus heterophyllus* (jak) leaves in streptozotocin-induced diabetic rats. Pharmacog mag 2010; 6(23):186-90.
5. Chandrika UG, Wedage WS, Wickramasinghe S. M. D. N and Fernando W. Hypoglycemic action of the flavanoid fraction of *Artocarpus heterophyllus* leaf. Afr. J. Trad. CAM 2006; 3(2):42-50.
6. Chiranjib Bhattacharjee and Amitsankar Dutta. Phytochemical and Acute Toxicity Study of Leaves of *Artocarpus Heterophyllus* Lam: International Journal of Allied Medical Sciences and Clinical Research (IJAMSCR), Volume 1, Issue 2, Nov – 2013
7. Dr C.K. Kokate, A.P. Parohit, S.B. Gokhale. Textbook of pharmacognosy: p:6.4-6.15
8. Dr. C.K Kokate. Practical textbook of pharmacognosy: p: 1-9.
9. Dr. C.K. Kokate, Dr D.S. Bidarks, S.B. Gokale. Textbook of pharmaceutical biotechnology: p:30-45.
10. Elevitch CR, Manner HI. *Artocarpus heterophyllus* (jackfruit). Species Profiles for Pacific Island agro forestry 2006; 1-17.
11. Gupta AK, Tandon N. Review on Indian Medicinal Plants. Indian Council of Medical Research: 2004.
12. Hakim Aliefman. Diversity of secondary metabolites from Genus *Artocarpus* (*Moraceae*). Nusantara Bio sci 2010; 2(3):146-156.
13. Jagtap UB, Bapat VA. *Artocarpus*: A review of its traditional uses, phytochemistry and pharmacology. J

Ethno pharmacol 2010; 129:142– 66.

14. Manjusha Choudhary, Vipin Kumar, Hitesh Malhotra,1 and Surender Singh. Medicinal plants with potential anti-arthritic activity: Journal of intercultural ethnopharmacology: 2015 Apr-Jun; 4(2): 147–179.
15. Om Prakash, Ritika Srivastava, Rajesh Kumar, Shivam Mishra, Shivam Srivastava. Preliminary Pharmacognostic and Phytochemical Studies on leaves of *Artocarpus heterophyllus*: International Journal of Natural Products and Marine Biology vol. 1 issue 1, April-2015; 35 – 40.
16. Patil KS, Jadhav AG, V. S Joshi. Wound healing activity of leaves of *Artocarpus heterophyllus*. Ind J Pharm Sci 2005; 67:629–32
17. Qiang Guo, Yuxiang Wang, Jiake Xu. Rheumatoid arthritis: Pathological mechanism and modern pharmacologic therapies: Bone research, 27 April 2018.
18. R A S N Ranasinghe, S D T Maduwanthi, R A U J Marapana. Nutritional and Health Benefits of Jackfruit (*Artocarpus heterophyllus Lam.*): A Review: Int J Food Sci 2019 Jan 6; 2019:4327183.
19. Reda Ben Mrid, Najat Bouchmaa Hassan Ainani, Rachid El Fatimy, Gabriel Malka, Loubna Mazini. Anti-rheumatoid drugs advancements: New insights into the molecular treatment of rheumatoid arthritis: Biomedicine and pharmacotherapy 151 (2022) 113126.
20. Soraya Riyanti, Clara Sunardi, Sana Nurul Falah. Isolation of Flavonoid from Jackfruit Leaves (*Artocarpus heterophyllus Lam.*) and its Antioxidant Activity: ResearchGate April 2013.

