IJNRD.ORG ISSN : 2456-4184
INTERNATIONAL JOURNAL OF NOVEL RESEARCH AND DEVELOPMENT (IJNRD) | IJNRD.ORG
An International Open Access, Peer-reviewed, Refereed Journal

Impact of Ginseng in Pharmaceutical Field: A Systematic Review

Ankita Jana[1], Srikanta Chandra[2]

[1]Assistant Professor,[2]Assistant professor

Department of Pharmaceutics

International Institute Of Pharmaceutical Science & Research,Kalyani,Nadia,India

Abstract: Ginseng is a well-known therapeutic plant which has been accepted by different countries like China, Korea, Iran, North America and Japan since decades because of its potent therapeutic activity. Different therapeutic uses of Ginseng plant can be found and is still the topic for their investigation and applications. Ginseng is a deciduous plant which belongs to Araliaceae family, the two significant types of ginseng plant are *Panax ginseng* and *Panax quinquefolius*, the roots of these plants have significant therapeutic activity. Several ginsenoids and non-saponin constituents have been chosen by the researchers for distinguishing their digestive and metabolomics activity. In the previous decades, fast improvement of biological innovations has encouraged the progression of ginseng examine from numerous points of view. Ginseng has been utilized as a dietary enhancement in a few nations everywhere in the world. Several studies have shown that Ginseng and their derivatives have a high pharmaceutical potential in the prevention and treatment of various diseases. Ginseng used as a dietary supplementary in the form of herbal tea, dried herbs, powder, and capsules. We believe that traditional herbs in the future open up new horizons for the pharmaceutical industry applications.

IndexTerms: Ginseng, Pharmacology of Ginseng, Chemoprevention, Health care by Ginseng

INTRODUCTION:

In USA ginseng plant consider among the main ten selling, home grown which is best used for dietary enhancement. E.g.*P.ginseng* which is also called red Korean ginseng & the uses of this plant mostly in the nourishment of chocolate, tea granular, nectar with cuts protected root, jam&confections (Leung and Wong 2010) [1-2]. In china also the ginseng plants which are having maturation span of five years are utilized as a potent nourishment material which significantly expanded the popularity of Ginseng plant (Leung and Wong 2010).The most popular Ginseng variety is the American ginseng (Panaxquinquefolius) and the Chinese ginseng (Panax notoginseng) which are widely used all over the world [3-6]. The polysaccharide components of ginseng plant has been considered due to their characteristic of developing various different organic exercises, such as antibacterial, radio protective, anti-mutagenic develop oxidation against ulcer, septicaemia and calming exercises (Sun Y, 2011).In the P. *ginseng* and p. *notoginseng* variety different polysaccharides have been found like arabinogalactan, gelatins and different acidic polysaccharides.They are comprised of a perplexing chain of monosaccharides wealthy in L-arabinose, D-galactose, L-rhamnose, D-galacturonic corrosive, D-glucuronic corrosive and D-galactosyl deposits (Wang et al., 2004). Among different chemical constituents of ginseng ginsenosides are the tri-terpene saponins having significant pharmacological action [7-10]. The skeleton of ginsenosides is made up of dammarane (four ring structure having 7 ring carbons) along with the sugar moieties (such as glucose, rhamnose, arabinose, xylose etc) attached at C-3 and C-20 positions. The ginsenosides are represented by the symbol Rx in which “R” represents root part of this plant and ”x” represents chromatographic extremity in an sequential order request (Leung and Wong 2010). Rb is the polar compound of this plant as compared to ra compound.

30 different varities of ginsenosides are characterized and divided into two classes the first one is twenty s protopanaxadiol which ranges from rb1 to rb3 other includes Rs1, Rh2, Rg3. Rd, Rc etc [11-15]. The second one is protopanaxatriol which includes re, rg2, rg, rf, rh1. The difference between the two classes is the gathering of carboxyl functional group at carbon 6 position in case of protopanaxadiol. A very few rare ginsenosides like 3,28-o-bisdesmoside) the pentacyclic saponin oleanane ro& f11 saponin ocotillol (r-24-pseudo-ginsenoside) (Leung and Wong 2010) are also found in the plant [16-20].Flawless and changed Ginsenosides are assimilated and moved to the human framework stays tricky.Transport of ginsenosides over the intestinal mucosa is vitality subordinate and non-saturable (Leung and Wong 2010).The plant contains different fixings, ginsenosides assume a more critical job in applying pharmacological activities than some other constituents [21-26 number of ginsenosides present in P. ginseng, less than 10 record for most ginsenoside substance. Specifically, ginsenosides rb1, rb2, rc, rd, re, rf, and rg1 are generally rich in the foundations of crude ginseng.Intriguingly, synthetic responses during the preparing of ginseng, for example, oxidation, hydrolysis, and additionally lack of hydration, lead to the arrangement of artifactual mixes, which regularly have upgraded natural exercises [27-30]. Plus, orally controlled ginsenosides experience bio-transformations in the gastrointestinal tract, and a few metabolites delivered by the activity of microorganisms have structures not the same as those of normally happening ginsenosides. Here, >100 ginsenosides have been arranged by their auxiliary Features (Byong-Kyu et al., 2015). In Proto-panaxadiols c3 & c20 are sugar moieties of ginsenosides which are dammarane type. In the terminal glucose of a 3-sugar chain the linkage happening b/w acylation & the chains of glycosyl at the 6-hydroxyl group [31-36]. Inside this gathering some structures are recognized include rb1 to rb3 & other including rh2, rg3, rd&rc. Almost thirty ginsenoside having a place with protopanaxadiol type which are characterised to the gathering of Rb. Proto-panaxatriols ginsenosides are dammarane type & they incorporate with ginsenosides Rf, Re, Rg1, Rg2, Rh1, F1 & F3 & not incorporate with R1 ginsenoside. In any case, in the protopanaxatriol moiety there is a hydroxyl branch at carbon 6 distinctive them from protopanaxadiol. Different highlights of protopanaxatriol structures are all things considered 2 chains of glycosyl & a straight linkage chains of saccharide (Yang et al.,2014). Rb1, Re, Rd, Rg1 & Rb3 ginsenosides are believed as 6 important saponins & comprise over seventy percent of ginsenoside absolute content in AG(Wang et al., 2005).

2

Oleanolic corrosive is another major t constituent, which have penta-cyclic triterpene skeleton (Shukla et al., 1997) [37-41]. Its secondary component is ginsenoside Ro (Huang et al., 2018). To wrap things up ocotillol that has a 5-membered epoxy ring at Carbon 20 should be referenced (Kim *et al*., 2012). From the leaves & roots of AG is pseudo-ginsenoside f11 detached ocotillol type panaxoside (Liu et al., 2017). Aside from polysaccharides & saponins AG contains A. A,phenolic mixes, terpenes, nutrients, flavonoids, unpredictable oils & minerals (Cui *et al*., 2017).An investigation conducted by Kochanet al. resulted that the constituent of triterpene saponins in shaggy root of AG can be distended by application of trans-anethole as elicitator (Kochanet al., 2018) [42-44].

Fig 1. Ginseng seeds

PHARMACOLOGICAL ACTIVITY OF GINSENG:

The ginseng plant Fig 1 is approved because of its wide range of pharmacological activities and large group of nearness to the bioactive cocktail. Ginsenosides are the essential ones including saponins steroidal which conjugated with different sugar polysaccharides & moieties which represent almost 10 to 20 percent of ginseng weight. Cell rein-forcement, calming and immunostimulatory exercises appear to be identified with the potential instruments of ginseng. The ginseng plant and its major constituent i.e. ginsenoside are clinically used for anti-diabetic, immune-modulatory, prophylactic, anti-cancer and different other activities on focal sensory system and for improving CVS framework [45-49]. The American ginsengs have immune-modulatory and prophylactic actionon respiratory sickness too (McElhaney et al., 2004; Predyet al., 2006; Vohra et al., 2008) Fig.2.

Fig 2. Pharmacological Activity of Ginseng

3

FOCAL SENSORY SYSTEM AND THEIR IMPACTS OF GINSENG:

The different neuro-degenerative disorders and other problems related to focal sensory system can be resolved using ginseng which basically imparts maturing impact (Christensen et al., 2009). ginsenosides shows significant job impact (Rausch et al., 2006). The defensive impacts of rb1 genocides (Chen et al., 2008), rg1 (Liu et al., 2010), rg3 (Tian and Fu, 2006), rd (Ye et al., 2009), and re (Chen et al., 2008c) on neuro-degeneration have been all around examined, in creatures & in neuronal societies of cell. When we study further than it shows that the sub-lingual vein infusion of the Ginsenosides rg3 at the portion of 5mg/kg displayed important neuro-protective impact on rodents for cerebral central ischemic damage related problem with decrease in the scores of neurological deficiency & reduced the territory infarct & done improvement in cerebral stream of blood [50]. Simultaneously, Rg3 could altogether improve mitochondrial vitality digestion, offend the declines in gsh,px& SOD exercises & increment in MDA level prompted by cerebral ischemia, proposing that Rg3 works through the decrease of peroxides lipids, rummaging free radicals & improving the vitality digestion (140005D).

GINSENG AND CARDIOVASCULAR:

The active components of ginseng can stimulate nitrogen oxide production, inhibit ROS production, increase blood circulation and help to adjust lipid profiles. In United States the American ginseng is a very popular domestic plant which enhances the health of people experiencing cardiovascular sickness (Xieet al., 2005a; Wang et al., 2007b). It has also shown anti-ischaemic, antiarrhythmic and antihypertensive impact (Wang et al., 2007b) [51]. The cancer prevention agent exercises & the connection b/w compound structure & cardiovascular-ensuring capacities have been evaluated (Prior and Cao, 2000; Wang et al., 2007b). American ginseng extricates had a more grounded cancer prevention agent movement than Asian ginseng root (Shao et al., 2004).

With the management of peroxide detoxifying systems the berry and ginseng root of American origin has shown a promising activity for preventing cancer and also defensive activity I refined (Mehendaleet al., 2006; Shao et al., 2004) and enacting the Nrf2 pathway (Li

et al., 2010b). Ginsenoside re was one of important cell reinforcement specialist that secured cardiomyocytes by searching hydrogen peroxide and hydroxyl radicals (Xieet al., 2006).

GINSENG AND ANTI-DIABETIC IMPACTS:

The most common type of diabetes is type 2 diabetes, which is a metabolic health problem with in appropriate digestion of starch and lipids due to weak insulin release (Qi et al., 2010a).Both American and Asian ginsengs show hypo-glycemic activity in the diabetic mouse model. (Atteleet al., 2002; Dey et al., 2003; Xieet al., 2002).

One of the important mouse modelob/ob is utilized; we showed that after twelve days of treatment with berry & ginseng American leaf decrease the level of fasting blood glucose & improve the transfer of glucose & reduced the weight of body (Atteleet al., 2002; Xieet al., 2004a; Xieet al., 2004b). When warmed American ginseng is compared with the natural ginseng the former one has shown more powerful impact in supressing impelled gathering of glycation end products in the kidney of diabetic mice (Kim et al., 2007a) [51]. The glucose level in plasma decreases and insulin secretion is increased by release of acetylcholine from the nerve terminals and binding with the muscarinic receptors in the cells of pancreas by ginsenoside rh2 (Lee et al., 2006). Through tweak of Akt/Foxo1/PDX-1 flagging pathway Rh2 has the potential to alter weakened cell development process and can supress apoptotic pathway (Wang et al., 2012). It was observed that comparitively Rg3 can improve work of islet cells and decreases the work of cytokine induced damage associated with nitric oxide creation and apoptosis in mouse (129). 25mg/kg oral dose of potassium could bring down the level of glucose plasma, triglycerides, cholesterol & level of NEFA by 20.7%, 41.6%, 20.2%, & 24.6%, separately (Yang et al., 2015). Ginseng is used as a traditional medicine for the treatment of DM in several countries such as China, Korea and Japan [52]. The ethanol extracts from wild gat-home significantly reduced blood sugar levels in fasting state according to dose used.

GINSENG AND CANCER CHEMOPREVENTION:

American ginseng & their important constituents show other pharmacological action in growth of malignant shows chemo-preventive & supress the development of tumour (Qi et al., 2010b; Wang et al., 2007); American ginseng separately updated the chemical and preventive effects of fluorouracil-5 (Li I et al, 2009d) in human colon cells, smoky chromosome variability initiated by mitomycin C in the mice (Pawar et al., 2007), improved the malignant depletion associated with plant growth (Barton et al., 2010), And provided radio potential in human lymphocytes (Lee et al., 2008b; Lee et al., 2010), (Wang et al., 2009a) [52].

GINSENG AND ANTI-CARCINOGENIC EFFECTS:

Ginsenosides is a major part of ginseng which when applied in vitro and in-vivo has shown promising anti-carcinogenic activity against few malignant growth types. Some of the ginsenosides have shown solid clinical outcomes and many others have been utilized against malignant growth items. The rough part of Ginseng's saponin applied more grounded in vitro cytotoxic exercises than ginseng red against ACFIN, 15-HCT & PC-3 cell lines with IC50 values running from 60.3–90.8 g/mL (Kim et al., 2008).

ANTI-INFLAMMATORY AND IMMUNE-MODULATORY EFFECTS OF GINSENG:

From a near report, dark Ginseng displayed more grounded calming and hostile to nociceptive impacts than ginseng red in instigated xylene ear oedema model in mice & carrageenan paw prompted oedema in rodents; it restrained the expert incendiary middle people, nitric oxide synthase inducible & 2 cyclo-oxygenase & professional cytokines fiery, interleukin 1 & 6 &tumour rot factor-(TNF-). What's more, the lipopolysaccharide-incited TNF-discharge was fundamentally diminished after treatment with dark ginseng remove (Lee et al., 2019). Besides, the part subordinate recovering impact of dark ginseng distinguish against the induced cisplatin nephron-toxicity & the decreased (pig cell LLC-PK1) cells reasonability, has been explained (Han et al., 2016) [53]. What's more, in a analogous test, compound potassium could annul the increased level of apoptotic LLC-PK1 cells altogether.

HEPATO-PROTECTIVE EFFECT OF GINSENG:

Ginseng dark has shown well hepatoprotective activity on acet-aminophen actuated mice the level of serum amino-transferase (AST) and lipid peroxidation alanine decreasing by liver damage compound malondialdehyde (MDA). In the meantime, the cell reinforcement levels were raised in the liver tissues in which include glutathione, p450 e1 cytochrome & hydroxynonenal 4(4-HNE). Treatment with ginseng dark also separate ethanol and decrease the gathering of lipid in the liver &harm the muscle of diabetic mice through the initiation of AMP-enacted protein kinase (AMPK) (Kang et al., 2017).

GINSENG AND DHOSTILE TO OBESITY AND EFFECTS OF ANTI-HYPERLIPIDAEMIC:

Ethanol ginseng concentrate have the potential to decrease hyperlipidaemia and also helps in the collection of the fat in fat regimen mice by the hindrance processing of fat. It has been seen that ginseng can efficiently remove as the same time it can decrease the level of complete cholesterol serum and low-density lipoprotein level in STZ-induced diabetic mice and fat eating mice (Saba et al., 2016). In male C57BLKS/J-db/db diabetic mice after ginseng treatment, a progressive decrease in triglyceride and unsaturated fat (NEFA)

levels was observed with expansion of the high-density level (Kim et al., 2012) [54]. H_2O and dark ginseng ethanol concentrates decrease lipid collection through the recommendation of phosphorylation PPAR, C/EBP and AMPK in 3T3-L1 cells with more anchored movement for ethanol extrication (Park et al., 2015).

TONIC EFFECT OF GINSENG:

Dark Ginseng medicines in a portion of 150 mg/kg have completely extended the rodent activity limit. The level of corrosive effect of lactic acid in the blood has decreased, but the movement of citrate synthase in the muscle has been extended (Jo and al, 2010). The dark form of ginseng may develop muscle development and may also treat muscle problems that are generally progressive and recognized by the production of myoblasts with large multinucleated myotubules , and increase breathing and thickness. This activity is considered to be apparent due to the initiation of the Act/mTOR/p70S6k concentrator (Lee et al., 2018).

TOPICAL USES OF GINSENG:

The dark matured ginseng has shown to play a significant negative impact on the wrinkle at 0.3g/ml convergence with the development of sort I pro-collagen articulation levels and reducing the MMP-1 articulation level. However, at 3 g/ml it significantly increased the outflow of TIMP-2 up to 154.55%. At 10 g/ml it decreases the articulation levels of both MMP-2 and MMP-9 to 45.15% and 66.65% (Pham et al., 2017).

Studies have demonstrated that Korean Red Ginseng (RG) shields skin from maturing and wrinkling, and it can likewise mitigate atopic dermatitis and sensitivity side effects. In a creature model, privately controlled ginseng radix rubra animated the inversion of diabetes and maturing in rodents (Morisakiet al., 1995) explores different avenues regarding rodents, affirming that ginseng expanded lipid and dampness content in skin. In the orally controlled RGp gathering, lipid and dampness content was higher than that in the benchmark group, and the distinctions were factually critical. Likewise, by contrasting injured skin tissue and typical skin tissue, we could reason that lipid and dampness substance and skin flexibility in injured tissue were practically identical with those in ordinary skin on postoperative day 20. Contrarily, there was no projective difference in skin lipid and dampness material or adaptability in the topically applied RGe bunch contrasted and those in the Vaseline gathering. The rapid effect that RG treatment had on wound mending is thought to have inflated the proportion of dampness. (Kee-sooet al., 2019).

GASTROINTESTINAL SYSTEMS:

AG isolates (fifty, hundred, & 200 mg/kg between) protected against oesophageal harm came about because of oesophagitis reflux by reduced of incendiary cells & oxidative burden in the throat (Singh et al., 2012). Another study exhibited which AG (500 and 1250 mg/kg between) shielded rodents from the mucosa of gastric harm caused by an incessant ethanol admission (Huang et al., 2013) [54].

GINSENG AND HOSTILE PROPERTIES OF AGEING:

For checking the impact of Ag by controlled clinical preliminaries were not yet performed. Many investigations were done by the researchers on the both ordered & natural parts so that they relate them for the prevention therapy of cancer agent characteristics of AG. 2.25g/kg for fourteen days of AG theorized to secure against untimely ovarian disappointment (POF), when the ovary's ability was lost before the age of 40, it will be considered as a marker for living maturation [55]. This review showed that AG provided against POF through the management of the prostaglandin biosynthesis, ovulation and anticipation of ovarian maturation (Zhu et al., 2012).

CONCLUSION:

The ginseng is used as traditional plant found in most countries of Asian for more than thousands of years. Ginseng contains a large number of active ingredients, as steroid mononides, protopanaksadiolki and protopanaksatriolki, collectively and called ginsenosides. This may be a valuable resource for the future development of drugs, but further evidence of higher quality natural product used in our life. In addition, ginseng may interact with medicines, although the evidence available suggests that this is a relatively safe product may be used in industry and pharmaceutical application, traditional medicine and food.

REFERENCES:

1)Attele AS, Zhou YP, Xie JT, Wu JA, Zhang L, et al. (2002). Antidiabetic effects of Panax ginseng berry extract and the identification of an effective component. Diabetes.; 51:1851–1858

2)Barton DL, Soori GS, Bauer BA, Sloan JA, Johnson PA, et al. (2010). Pilot study of Panax quinquefolius (American ginseng) to improve cancer-related fatigue: a randomized, double-blind, dose-finding evaluation: NCCTG trial N03CA. Support Care Cancer. 18:179–187

3)Byong-Kyu Shin, Sung Won Kwon, Jeong Hill Park. (2015). Chemical diversity of ginseng saponins from Panax ginseng. J Ginseng Res 39 :287e298

4)Chen JX, Peng HM, Xi OY, He XY. (2008 b). Research on the antitumor effect of ginsenoside Rg3 in B16 melanoma cells. Melanoma Res. 18:322–329.

5)Christensen LP. (2009). Ginsenosides chemistry, biosynthesis, analysis, and potential health effects. Adv Food Nutr Res. 55:1–99.

6)Cui S.;Wu J.;Wang J.;Wang X. (2017). Discrimination of American ginseng and Asian ginseng using electronic nose and gas chromatography mass spectrometry coupled with chemometrics. J. Ginseng Res. 41, 85–95.

7)Dey L, Xie JT, Wang A, Wu J, Maleckar SA, Yuan CS. (2003). Anti-hyperglycaemic effects of ginseng: comparison between root and berry. Phytomedicine. 10:600–605.

8)Han M.-S.; Han I.-H.; Lee D.; An J.M.; Kim S.-N.; et al. (2016). Beneficial effects of fermented black ginseng and its ginsenoside 20 (S)-Rg3 against cisplatin- induced nephrotoxicity in LLC-PK1 cells. J. Ginseng Res. 40, 135–140.

9)Huang C.C.; Chen Y.M.; Wang D.C.; Chiu C.C.; Lin W.T.; et al. (2013). Cytoprotective effect of American ginseng in a rat ethanol gastric ulcer model. Molecules, 19, 316–326

10)Huang X.; Liu Y.; Zhang N.; Sun X.; Yue H.; et al. (2018). UPLC Orbitrap HRMS analysis of Panax quinquefolium L. for authentication of Panax genus with chemometric methods. J. Chromatogr. Sci. 56, 25–35.

11)Jo G.S.; Chai H.-Y.; Ji H.J.; Kang M.H.; Kang S.-J.; et al. (2010). Enhancement of exercise capacity by black ginseng extract in rats. Lab. Anim. Res. 26, 279–286.

12)Kang, O.-H.; Shon, M.-Y.; Kong, R.; Seo, Y.-S.; Zhou, T.; et al. (2017). Anti-diabetic effect of black ginseng extract by augmentation of AMPK protein activity and upregulation of GLUT2 and GLUT4 expression in db/db mice. Bmc Complementary Altern. Med., 17, 341

13)Kar Wah Leung and Alice Sze-Tsai Wong. (2010). pharmacology of ginsenosides: a literature review. Chinese Medicine, 5:20

14)Kim SY, Kim DH, Han SJ, Hyun JW, Kim HS. (2007b). Repression of matrix metalloproteinase gene expression by ginsenoside Rh2 in human astroglioma cells. BiochemPharmacol. 74:1642–1651

15)Kim, A.-J.; Kang, S.-J.; Lee, K.-H.; Lee, M.; Ha, S.-D.; et al. (2010). The chemo preventive potential and anti-inflammatory activities of Korean black ginseng in colon26-M3. 1 carcinoma cells and macrophages. J. Korean Soc. Appl. Biol. Chem. 53, 101–105.

16)Kim, A.-J.; Yoo, H.-S.; Kang, S.-J. (2012). Ameliorative effect of black ginseng on diabetic complications in C57BLKS/J-db/db mice. Korean J. Food Nutr. 25, 99–104.

17)Kim, D.H. (2012). Chemical diversity of Panax ginseng, Panax quinquefolium, and Panax notoginseng. J Ginseng Res., 36, 1–15.

18)Kim, E.-K.; Lee, J.-H.; Cho, S.-H.; Shen, G.-N.; Jin, L.-G.; et al. (2008). Preparation of black panax ginseng by new methods and its antitumor activity. Korea J. Herbol. 23, 85–92.

19)Kim, S.; Jang, H.; Oh, M.; Eom, D.; Kang, K.; et al. (2014). Ginsenoside Rg3 enhances islet cell function and attenuates apoptosis in mouse islets. In Transplantation Proceedings; Elsevier: Amsterdam, The Netherlands, Volume 46, pp. 1150–1155.

20)Ki-Soo Park, Dae-Hwan Park. (2019). The effect of Korean Red Ginseng on full-thickness skin wound healing in rats. J Ginseng Res 43: 226e235

21)Kochan, E.; Szymczyk, P.; Ku´zma, Ł.; Szyma´ nska, G.; Wajs-Bonikowska, A.; et al. (2018). The increase of triterpene saponin production induced by trans-anethole in hairy root cultures of Panax quinquefolium. Molecules, 23, 2674.

22)Lee LS, Wise SD, Chan C, Parsons TL, Flexner C, Lietman PS. et al. (2008a). Possible differential induction of phase 2 enzyme and antioxidant pathways by American ginseng, Panax quinquefolius. J Clin Pharmacol. 48:599–609

23)Lee TK, O'Brien KF, Wang WD, Johnke RM, Sheng C, et al. (2010). Radioprotective effect of American ginseng on human lymphocytes at 90 minutes postirradiation: A study of 40 cases. J Altern Complement Med. 16:561–567.

24)Lee, S.-Y.; Go, G.-Y.; Vuong, T.A.; Kim, J.W.; Lee, S.; et al. (2018). Black ginseng activates Akt signaling, thereby enhancing myoblast differentiation and myotube growth. J. Ginseng Res. 42, 116–121.

25)Lee, W.K.; Kao, S.T.; Liu, I.M.; Cheng, J.T. (2006). Increase of insulin secretion by ginsenoside Rh2 to lower plasma glucose in Wistar rats. Clin. Exp. Pharmacol. Physiol., 33, 27–32.

26)Lee, Y.Y.; Saba, E.; Irfan, M.; Kim, M.; Yi-Le Chan, J et al. (2019). The anti- inflammatory and anti-nociceptive effects of Korean black ginseng. Phytomedicine. 54, 169–181.1227.

27)Li HF, Ye M, Guo HZ, Tian Y, Zhang J, et al. (2009b). Biotransformation of 20(S)- protopanaxadiol by Mucor spinosus. Phytochemistry. 70:1416–1420.

28)Liu, Y.Y.; Zhang, T.Y.; Xue, X.; Liu, D.M.; Zhang, H.T.; et al. (2017). Pseudoginsenoside-F11 attenuates cerebral ischemic injury by alleviating autophagic/lysosomal defects. CNS Neurosci. Ther., 23, 567–579.

29)McElhaney JE, Gravenstein S, Cole SK, Davidson E, O'Neill D, et al. (2004). A placebo-controlled trial of a proprietary extract of North American ginseng (CVT-E002) to prevent acute respiratory illness in institutionalized older adults. J Am Geriatr Soc. 52:13–19.

30)Mehendale SR, Wang CZ, Shao ZH, Li CQ, Xie JT, et al. (2006). Chronic pretreatment with American ginseng berry and its polyphenolic constituents attenuate oxidant stress in cardiomyocytes. Eur J Pharmacol. 553:209–214

31)Morisaki N, Watanabe S, Tezuka M, Zenibayashi M, Shiina R, et al. (1995). Mechanism of angiogenic effects of saponin from ginseng radix rubra in human umbilical vein endothelial cells. Br J Pharmacol;115:1188e93.

32)Park, H.-J.; Kim, A.-J.; Cheon, Y.-P.; Lee, M. (2015). Anti-obesity effects of water and ethanol extracts of black ginseng. J. Korean Soc. Food Sci. Nutr. 44, 314–323

33)Pawar AA, Tripathi DN, Ramarao P, Jena G. (2007). Protective effects of American ginseng (Panax quinquefolium) against mitomycin C induced micronuclei in mice. Phytother Res.; 21:1221–

34)Pham, Q.L.; Jang, H.J.; Kim, K.B. (2017). Anti-wrinkle effect of fermented black ginseng on human fibroblasts. Int. J. Mol. Med., 39, 681–686

35)Predy GN, Goel V, Lovlin RE, Basu TK. et al. (2006). Immune modulating effects of daily supplementation of COLD-fX (a proprietary extract of north American ginseng) in healthy adults. J Clin BiochemNutr. 39:160–165.

36)Prior RL, Cao G. et al. (2000). Analysis of botanicals and dietary supplements for antioxidant capacity: a review. J AOAC Int. 83:950–956.

37)Qi LW, Wang CZ, Yuan CS. (2010b). American ginseng: Potential structure-function relationship in cancer chemoprevention. BiochemPharmacol. 80:947–954.

38)Rausch WD, Liu S, Gille G, Radad K. (2006). Neuroprotective effects of ginsenosides. Acta Neurobiol Exp.; 66:369–375.

39)Saba, E.;Jeon, B.R.; Jeong, D.-H.; Lee, K.; Goo, Y.-K.; et al. (2016). Black ginseng extract ameliorates hypercholesterolemia in rats. J. Ginseng Res. 40, 160–168.

40)Shao ZH, Xie JT, Vanden Hoek TL, Mehendale S, Aung H, et al. (2004). Antioxidant effects of American ginseng berry extract in cardiomyocytes exposed to acute oxidant stress. BiochimBiophys Acta. 1670:165–171.

41)Shukla,Y.N.; Tripathi, A.K.; Mehta,V.K. (1997). Feeding-deterrency of oleanolic acid isolated fromPanax quinquefolium against lepidopterans. Phytother. Res. 11, 591–593.

42)Singh, P.; Singh, N.; Sengupta, S.; Palit, G. (2012). Ameliorative effects of Panax quinquefolium on experimentally induced reflux oesophagitis in rats. Indian J. Med. Res. 135, 407–413

43)Sun Y (2011). Structure and Biological Activities of the Polysaccharides from the Leaves, Roots and Fruits of Panax ginseng C.A. Meyer: An Overview. Carbohyd. polym. 85: 490-499.

44)Tian J, Fu F. (2006). Neuroprotective effects of 20(S)-ginsenoside Rg3 on dopamine levels in the 6- hydroxydopamine lesioned mouse brain. Movement Disord.

45)Van Kampen J, Robertson H, Hagg T, Drobitch R. et al. (2003). Neuroprotective actions of the ginseng extract G115 in two rodent models of Parkinson's disease. Exp Neurol. 184:521–529.

46)Wang CZ, Aung HH, Ni M, Wu JA, Tong RB, et al. (2007a). Red American ginseng: Ginsenoside constituents and antiproliferative activities of heat-processed Panax quinquefolius roots. Planta Med. 73:669–674

47)Wang M, Guilbert LJ, Li J, Wu Y, Pang P, et al.(2004) A Proprietary Extract from North American Ginseng (Panax quinquefolium) Enhances IL-2 and IFN-gamma Productions in Murine Spleen Cells Induced by Con-A. Int. immunopharmacol. 4:311–315.

48)Wang, A.B.; Wang, C.Z.; Wu, J.A.; Osinski, J.; Yuan, C.S. et al. (2005). Determination of major ginsenosides in Panax quinquefolius (American ginseng) using high-performance liquid chromatography. Phytochem. Anal. 16, 272–277.

49)Wang, Y.;Wang, H.; Liu, Y.; Li, C.; Qi, P.; Bao, J. et al. (2012). Antihyperglycemic effect of ginsenoside Rh2 by inducing islet -cell regeneration in mice. Horm. Metab. Res. 44, 33–40.

50)Xie JT, Shao ZH, Vanden Hoek TL, Chang WT, Li J, et al. (2006). Antioxidant effects of ginsenoside Re in cardiomyocytes. Eur J Pharmacol. 532:201–207

51)Xie JT, Wu JA, Mehendale S, Aung HH, Yuan CS. et al. (2004b). Anti-hyperglycemic effect of the polysaccharides fraction from American ginseng berry extract in ob/ob mice. Phytomedicine. 11:182–187.

52)Xie JT, Zhou YP, Dey L, Attele AS, Wu JA, et al. (2002). Ginseng berry reduces blood glucose and body weight in db/db mice. Phytomedicine. 9:254–258.

53)Yang, X.-D.; Yang, Y.-Y.; Ouyang, D.-S.; Yang, G.-P. et al. (2015). A review of biotransformation and pharmacology of ginsenoside compound K. Fitoterapia, 100, 208–220

54)Yang,W.Z.; Hu, Y.;Wu,W.Y.; Ye, M.; Guo, D.A. (2014). Saponins in the genus Panax L. (Araliaceae): A systematic review of their chemical diversity. Phytochemistry, 106, 7–24.

55)Zhu, L.; Li, J.; Xing, N.; Han, D.; Kuang, H.; Ge, P. (2015). American ginseng regulates gene expression to protect against premature ovarian failure in rats. Biomed Res. Int., 2015, 767124.